

干枝病治疗法

Rawatan Penyakit Pucuk Kering Twig Blight Treatment



INOGuard

+



SNP



CGKing

+



SBIO-C



INOGuard	SNP	Water 水
1L	4L	1000L
200ml	800ml	200L
20ml	80ml	20L

治疗法 Rawatan Treatment	• INOGuard + SNP + 水, 喷洒整棵树 • 每7-10天喷洒一次, 至少三次 • 每两星期施肥 SBIOC + CGKing 补足营养	• INOGuard + SNP + Air, sembur pada seluruh pokok • Semburkan dalam setiap 7-10 hari, minima 3 kali • Tabur baja SBIOC + CGKing setiap 2 minggu	• INOGuard + SNP + Water, spray onto entire tree • Spray every 7-10 days, at least 3 times • Apply SBIOC + CGK every 2 weeks
预防法 Pencegahan Prevention	• INOGuard + SNP + 水, 喷洒整棵树 • 每月喷洒一次 • 施肥工作如常	• INOGuard + SNP + Air, sembur pada seluruh pokok • Semburkan dalam setiap bulan • Tabur baja seperti biasa	• INOGuard + SNP + Water, spray onto entire tree • Spray every month • Apply fertilizer as usual

注：喷叶底为佳，适于上午十点前或下午五点后喷

Nota: Galakkan penyemburan foliar pada bawah daun pada waktu sebelum 10am atau selepas 5pm

Note: Encourage spray onto underneath leaf, spray before 10am or after 5pm